



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 429 281 A2

12

EUROPEAN PATENT APPLICATION

21 Application number: 90312579.7

51 Int. Cl.⁵: C07C 279/18, A01N 47/44

22 Date of filing: 19.11.90

30 Priority: 20.11.89 JP 299655/89
04.10.90 JP 265230/90

43 Date of publication of application:
29.05.91 Bulletin 91/22

84 Designated Contracting States:
DE FR GB IT

71 Applicant: HOKKO CHEMICAL INDUSTRY CO.,
LTD
4-20, 4-chome, Nihonbashi-Hongoku-cho
Chuo-ku
Tokyo(JP)

72 Inventor: Ishikawa, Hiromichi
40-12, Asahi-cho 5-chome
Atsugi-shi, Kanagawa-ken(JP)
Inventor: Umeda, Takashi
5-2-306, Sagamidaidanchi


Sagamihara-shi, Kanagawa-ken(JP)
Inventor: Onoue, Shinji c/o Hokko Kagaku-Ryo
2385, Toda

Atsugi-shi, Kanagawa-ken(JP)
Inventor: Kajikawa, Kazuo
c/o Hokko Kagaku-Ryo, 2385, Toda
Atsugi-shi, Kanagawa-ken(JP)
Inventor: Shibata, Toshihiro
c/o Hokko Kagaku-Ryo, 2385, Toda
Atsugi-shi, Kanagawa-ken(JP)
Inventor: Ohyama, Hiroshi
B-22-19, 36-20, Tsutsumi
Chigasaki-shi, Kanagawa-ken(JP)

74 Representative: Cresswell, Thomas Anthony
et al
J.A. Kemp & Co. 14 South Square Gray's Inn
London WC1R 5LX(GB)

54 Guanidine derivatives and fungicides for agriculture and horticulture containing the same.

57 The guanidines of formula (I)



wherein R is hydrogen, (C₁-C₁₀)alkyl, (C₂-C₆)alkenyl, (C₂-C₄)-alkynyl, (C₅-C₆)cycloalkyl, (C₁-C₆)alkyl or (C₅-C₇)-cycloalkyl optionally substituted with (C₁-C₆)alkyl, X is hydrogen, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy, Y is hydrogen, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy, R¹ and R² are the same or different and each is hydrogen or methyl, and l, m and n are the same or different and each is an integer of 1 or 2 provided that when R, X, Y, R¹ and R² are all hydrogen, l is 2, and acid addition salts thereof possess high control effects on mildew, blight, powdery mildew and rust which are serious diseases of fruit trees, vegetables or cereals, and are useful as agricultural and horticultural fungicides.

EP 0 429 281 A2

GUANIDINE DERIVATIVES AND FUNGICIDES FOR AGRICULTURE AND HORTICULTURE CONTAINING THE SAME


Background of the Invention

1. Field of the Invention

This invention relates to novel guanidine derivatives and fungicides for agriculture and horticulture containing said derivatives as an active ingredient.


2. Description of the Prior Art

Some of the literatures disclose guanidines. For example, Japanese Patent Publication No. 29742/1968 discloses a process for preparing substituted guanidines represented by the formula



wherein R_1' represents a substituted or unsubstituted aromatic hydrocarbon residue and R_2' or R_3' represents a hydrogen atom, a substituted or unsubstituted aromatic hydrocarbon residue.

There are also disclosed in Japanese Patent Publication No. 9846/1975 guanidines represented by the formula




wherein X_1' represents a hydrogen atom or a chlorine atom and R_4' represents a benzylamino group or the like which have insecticidal and acaricidal activities as well as plant disease controlling activity.

Although the known guanidines have some fungicidal activities for agriculture and horticulture, they have not been put into practical use due to their not satisfactorily high activities and their phytotoxicities. On the other hand, serious diseases of fruit trees, vegetables or cereals such as mildew, blight, powdery mildew and rust have hitherto been treated with various chemicals, all of which are unusable or of restricted use due to resistance to the chemicals. Therefore, development of novel fungicides different in skelton structure from prior-art chemicals has become a big problem in these fields.

Detailed Description of the Invention

To attain the above-mentioned object, we have synthesized a large number of compounds and studied their usefulness. As a result, we were successful in the synthesis of the guanidine derivatives of the below-mentioned formula (I), and found that these derivatives are new compounds not disclosed in any literature and furthermore they have higher fungicidal activity and safety for agriculture and horticulture.

Therefore, the first aspect of the invention is guanidine derivatives represented by the formula



wherein R represents a hydrogen atom, a (C₁-C₁₀)-alkyl group, a lower alkenyl group, a lower alkynyl group, a (C₅-C₆)-cycloalkyl-lower alkyl group or a (C₅-C₇)-cycloalkyl group which may be substituted with a lower alkyl group, X represents a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy group, a lower haloalkyl group or a lower haloalkoxy group, Y represents a hydrogen atom, a halogen atom, a lower alkyl group or a lower alkoxy group, R₁ and R₂ independently represent a hydrogen atom or a methyl group, and t, m and n independently represent an integer of 1 or 2 except that R, X, Y, R₁ and R₂ each are a hydrogen atom and t is 1.

The lower alkyl group as used herein means an alkyl group having 1-6 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, t-butyl, pentyl or hexyl.

When R is a lower alkenyl group, it is an alkenyl group having 2-6 carbon atoms such as vinyl, allyl, 1-propen-2-yl, butenyl, pentenyl or hexenyl.

When R is a lower alkynyl group, it is an alkynyl group having 2-4 carbon atoms such as ethynyl, propargyl, 1-propyn-2-yl or butynyl.

When X and Y independently are a lower alkoxy group, they are an alkoxy group having 1-5 carbon atoms such as methoxy, ethoxy, propoxy, isopropoxy, butoxy or pentyloxy.

Examples of a cycloalkyl-lower alkyl group for R include cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl and cyclohexylethyl.


Examples of a (C₅-C₇)-cycloalkyl group optionally substituted with a lower alkyl group include cyclopentyl, cyclohexyl, cycloheptyl which are optionally substituted with methyl, ethyl, n-propyl, iso-propyl or n-butyl.

The second aspect of the invention is a fungicide for agriculture and horticulture containing as an active ingredient a dibenzylguanidine derivative of the above-mentioned formula (I).

Examples of the inventive compounds of the formula (I) are shown in Table 1.

It is to be noted that the compound Nos. will be referred to also in Examples and Test Examples below.

Table 1



Compound No.	R	X _m	Y _n	$\left(\begin{array}{c} \text{R}_1 \\ \\ \text{C} \\ \\ \text{R}_2 \end{array} \right)$	Physical data
1	H-	4-C/	H	CH ₂	n_D^{23} 1.6259
2	H-	4-CH ₃ -	H	CH ₂	n_D^{23} 1.6112
3	H-	2-CH ₃ O-	H	CH ₂	n_D^{23} 1.6308
4	CH ₃ -	H-	H	CH ₂	n_D^{23} 1.6071
5	CH ₃ -	2-C/-	H	CH ₂	m.p. 81-84°C
6	CH ₃ -	4-C/-	H	CH ₂	n_D^{23} 1.6042
7	CH ₃ -	4-CH ₃ -	H	CH ₂	n_D^{23} 1.5985
8	CH ₃ -	4-CF ₃ -	H	CH ₂	n_D^{23} 1.5994
9	C ₂ H ₅ -	H-	H	CH ₂	n_D^{23} 1.5966
10	C ₂ H ₅ -	4-C/	H	CH ₂	m.p. 73-75°C
11	n-C ₃ H ₇ -	H-	H	CH ₂	n_D^{23} 1.5831
12	n-C ₃ H ₇ -	3-C/	H	CH ₂	n_D^{23} 1.5884
13	iso-C ₃ H ₇ -	H-	H	CH ₂	n_D^{23} 1.5847
14	iso-C ₃ H ₇ -	4-C/-	H	CH ₂	m.p. 122-125°C
15	iso-C ₃ H ₇ -	4-tert-C ₄ H ₉ -	H	CH ₂	m.p. 96-97°C
16	n-C ₄ H ₉ -	2-CH ₃ -	H	CH ₂	n_D^{23} 1.6005

Table 1 (cont'd)










Com- pound No.	R	X _m	Y _n	$\begin{pmatrix} R_1 \\ \\ C \\ \\ R_2 \end{pmatrix}$	Physical data
17		H-	H	CH ₂	m.p. 106-107°C
18		2-C/-	H	CH ₂	m.p. 108-111°C
19		4-C/-	H	CH ₂	m.p. 140-141°C
20		2-CH ₃ -	H	CH ₂	n _D ²³ 1.6049
21		4-CHF ₂ O-	H	CH ₂	m.p. 86.5-88°C
22		H-	H	CH ₂	m.p. 164-166°C
23		4-C/-	H	CH ₂	m.p. 174-175°C
24		3-iso-C ₃ H ₇ O-	H	CH ₂	n _D ²³ 1.5946
25		4-CHF ₂ O-	H	CH ₂	m.p. 118-120°C
26	CH ₃ -	4-F-	H	CH ₂	n _D ²³ 1.6130
27	CH ₃ -	3-C/-	H	CH ₂	n _D ²³ 1.5842
28	CH ₃ -	4-CH ₃ O-	H	CH ₂	n _D ²³ 1.5555
29	iso-C ₃ H ₇ -	2-C/-	H	CH ₂	n _D ²³ 1.5470
30	iso-C ₃ H ₇ -	3-C/-	H	CH ₂	n _D ²³ 1.5571
31	iso-C ₃ H ₇ -	4-CH ₃ O-	H	CH ₂	m.p. 145-146°C
32	n-C ₄ H ₉ -	2-C/-	H	CH ₂	n _D ²³ 1.5269
33	sec-C ₄ H ₉ -	4-C/-	H	CH ₂	m.p. 120-122°C

Table 1 (cont'd)
















Com- pound No.	R	X _m	Y _n	$\begin{pmatrix} R_1 \\ \\ C \\ \\ R_2 \end{pmatrix}$	Physical data
34		2-F-	H	CH ₂	m.p. 126-127°C
35		3-C/-	H	CH ₂	m.p. 90-92°C
36		4-Br-	H	CH ₂	m.p. 159-160°C
37		4-CH ₃ -	H	CH ₂	m.p. 118-119°C
38		4-tert-C ₄ H ₉ -	H	CH ₂	m.p. 105-108°C
39		4-CH ₃ O-	H	CH ₂	m.p. 143.5-144.5°C
40		4-F-	H	CH ₂	m.p. 183-184°C
41		2-C/-	H	CH ₂	m.p. 147-148°C
42		3-C/-	H	CH ₂	m.p. 135-136°C
43		4-Br-	H	CH ₂	m.p. 164-172°C
44		2-CH ₃ -	H	CH ₂	m.p. 72-75°C
45		3-CH ₃ -	H	CH ₂	m.p. 128-129°C
46		4-CH ₃ -	H	CH ₂	m.p. 172-173°C
47		4-tert- C ₄ H ₉ -	H	CH ₂	m.p. 136-138°C
48		4-CH ₃ O-	H	CH ₂	m.p. 123-124°C
49	H-	2-C/-	4-CH ₃ -	CH ₂	n_D^{23} 1.6122
50	H-	3-C/-	4-CH ₃ -	CH ₂	n_D^{23} 1.6039

Table 1 (cont'd)

Compound No.	R	X _m	Y _n	$\begin{array}{c} \text{R}_1 \\ \\ \text{--- C ---} \\ \\ \text{R}_2 \end{array}$	Physical data
51	H-	4-C/-	4-CH ₃ -	CH ₂	n _D ²³ 1.6048
52	CH ₃ -	4-C/-	4-F-	CH ₂	n _D ²³ 1.5867
53	CH ₃ -	4-C/-	4-C/-	CH ₂	n _D ²³ 1.5354
54	CH ₃ -	4-C/-	4-CH ₃ -	CH ₂	n _D ²³ 1.5258
55	CH ₃ -	4-C/-	4-CH ₃ -	CH ₂	n _D ²³ 1.5786
56	CH ₃ -	4-C/-	2,6-(CH ₃) ₂ -	CH ₂	n _D ²³ 1.5603
57	CH ₃ -	4-CF ₃ -	3-C/-	CH ₂	n _D ²³ 1.5876
58	CH ₃ -	3-iso-C ₃ H ₇ -	2-CH ₃ -	CH ₂	n _D ²³ 1.5334
59	C ₂ H ₅ -	3-C/-	2-CH ₃ -	CH ₂	n _D ²³ 1.5842
60	C ₂ H ₅ -	4-C/-	2,6-(CH ₃) ₂ -	CH ₂	n _D ²³ 1.5729
61	C ₂ H ₅ -	2-CH ₃ -	2-C/-	CH ₂	n _D ²³ 1.5858
62	n-C ₃ H ₇ -	2-C/-	3-CH ₃ -	CH ₂	n _D ²³ 1.5702
63	n-C ₃ H ₇ -	4-CH ₃ -	3-C/-	CH ₂	n _D ²³ 1.5619
64	iso-C ₃ H ₇ -	2-C/-	2,4-C/2-	CH ₂	n _D ²³ 1.5375
65	iso-C ₃ H ₇ -	4-C/-	4-CH ₃ O-	CH ₂	n _D ²³ 1.5778
66	iso-C ₃ H ₇ -	4-C/-	2,6-(CH ₃) ₂ -	CH ₂	m.p. 107-110°C
67	iso-C ₃ H ₇ -	4-CF ₃ -	4-C/-	CH ₂	n _D ²³ 1.5444

Table 1 (cont'd)
















Com- pound No.	R	X _m	Y _n	$\begin{pmatrix} R_1 \\ \\ C \\ \\ R_2 \end{pmatrix}$	Physical data
68	n-C ₄ H ₉ -	4-C/-	3-CH ₃ -	CH ₂	n _D ²³ 1.5287
69		2-CH ₃ -	3-iso- C ₃ H ₇ O-	CH ₂	n _D ²³ 1.5763
70		3-CH ₃ -	2-CH ₃ -	CH ₂	n _D ²³ 1.5311
71		4-tert- C ₄ H ₉ -	2-C/-	CH ₂	n _D ²³ 1.5367
72		2-CH ₃ -	4-C/-	CH ₂	n _D ²³ 1.5803
73		4-CHF ₂ O-	4-CH ₃ -	CH ₂	n _D ²³ 1.5226
74		H-	4-F-	CH ₂	m.p. 173-175°C
75		H-	4-C/-	CH ₂	m.p. 145-147°C
76		H-	4-CH ₃ -	CH ₂	m.p. 128-130°C
77		H-	4-CH ₃ O-	CH ₂	m.p. 168-170°C
78		4-C/-	4-F-	CH ₂	m.p. 179-181°C
79		4-C/-	4-C/-	CH ₂	m.p. 172-174°C
80		4-C/-	4-CH ₃ -	CH ₂	m.p. 135-137°C
81		4-C/-	4-CH ₃ O-	CH ₂	m.p. 187-189.5°C
82		4-C/-	2,6-(CH ₃) ₂ -	CH ₂	m.p. 168-170°C
83		4-CF ₃ -	3-iso-C ₃ H ₇ O-	CH ₂	n _D ²³ 1.5952
84	H-	2,4-C ₂ -	H	CH ₂	m.p. 61-63°C

Table 1 (cont'd)











Compound No.	R	X _m	Y _n	$\begin{pmatrix} R_1 \\ \\ -C- \\ \\ R_2 \end{pmatrix}$	Physical data
85	CH ₃	2,4-C ₂ -	H	CH ₂	n _D ²³ 1.5749
86	iso-C ₃ H ₇	2,4-C ₂ -	H	CH ₂	m.p. 163-166°C
87		2,4-C ₂ -	H	CH ₂	m.p. 121-123°C
88		2,4-C ₂ -	H	CH ₂	m.p. 175-176°C
89	n-C ₁₀ H ₂₁ -	4-C/-	H	CH ₂	n _D ²³ 1.4914
90	CH ₂ =CHCH ₂ -	4-C/-	H	CH ₂	n _D ²³ 1.5466
91	CH≡CCH ₂ -	4-C/-	H	CH ₂	n _D ²³ 1.5526
92	CH ₃ CH=CHCH ₂ -	4-C/-	H	CH ₂	n _D ²³ 1.5497
93		4-C/-	H	CH ₂	m.p. 118-131°C
94		4-C/-	H	CH ₂	m.p. 98-100°C
95		4-C/-	H	CH ₂	m.p. 193-196°C
96		4-C/-	H	CH ₂	m.p. 162-164°C
97		4-C/-	H	CH ₂	n _D ²³ 1.5672
98		4-C/-	H	CH ₂	m.p. 131-132°C
99	H-	H-	H	$\begin{pmatrix} CH_3 \\ \\ -CH- \end{pmatrix}$	n _D ²³ 1.5022



Table 1 (cont'd)

Compound No.	R	X _m	Y _n	$\left(\begin{array}{c} R_1 \\ \\ -C- \\ \\ R_2 \end{array} \right)_l$	Physical data
100	H-	H-	H	$\begin{array}{c} CH_3 \\ \\ -C- \\ \\ CH_3 \end{array}$	n _D ²³ 1.4983
101	H-	H-	H	-CH ₂ CH ₂ -	n _D ²³ 1.5255
102		H-	H	-CH ₂ CH ₂ -	m.p. 108-109°C
103		4-C/-	H	-CH ₂ CH ₂ -	n _D ²³ 1.4924

The present compounds of the formula (I) are novel compounds which are effective as an active ingredient in fungicides for agriculture and horticulture.

Process for preparing the present compounds

The compounds of the formula (I) according to the invention can be prepared by the procedures as described below. The compounds can be prepared by reacting a phenylisocyanide dichloride represented by the formula (II) with a benzylamine derivative represented by the formula (III).



wherein R, X, Y, R₁, R₂, l, m and n have the same meanings as defined above.

The condensation reaction is usually carried out in an organic solvent. The solvent that may be

employed includes hydrocarbons such as toluene and hexane, halogenated hydrocarbons such as chloroform and chlorobenzene, ethers such as ethyl ether, dioxane and tetrahydrofuran, nitriles such as acetonitrile and propionitrile, alcohols such as methanol and ethanol and dimethylsulfoxide.

As the compounds of the formula (III) are basic substances, they may be used in excess in place of the acid-binding agent. Alternatively, inorganic bases such as sodium hydride, sodium amide, sodium hydroxide and potassium carbonate, and organic bases such as triethylamine and pyridine may be used as the acid-binding agent.

Whereas the reaction proceeds at room temperature, heating at a temperature up to boiling point of the solvent used can shorten the reaction time. After completion of the reaction, salts of the acid-binding agent, if any, are separated by filtration, and the desired product can be obtained by removing the solvent by distillation. Alternatively, the desired product can be separated by the addition of water and an organic solvent such as benzene, toluene, tetrahydrofuran or chloroform. The compound of the invention is obtained by removal of the solvent by distillation.

The starting compounds (II) and (III) are known compounds. Examples 1-4 specifically illustrate the preparation according to the above procedures.

Preparative Example 1 Preparation of 1,3-dibenzyl-1,3-diisopropyl-2-phenylguanidine (Compound No. 13)

In a 500-ml four-necked flask were placed 59.6 g of N-isopropylbenzylamine and 200 ml of acetonitrile. After cooling with water 17.4 g of phenylisocyanide dichloride was added dropwise. The mixture was then stirred at 50 °C for 3 hours. After cooling, precipitated salts were separated by filtration, the filtrate concentrated and finally water and toluene added to the residue. The toluene layer was washed with an aqueous 1N-hydrochloric acid and concentrated under reduced pressure to afford 36.7 g of the title compound as a pale yellow oil. The oil was purified by column chromatography on silica gel using a hexane-acetone mixture to give a colorless oil (yield 24.5 g), $n_D^{23} = 1.5847$.

Preparative Example 2 Preparation of 1,3-dicyclopentyl-1,3-di(4-difluoromethoxybenzyl)-2-phenylguanidine (Compound No. 21)

In a 500-ml four-necked flask were placed 48.2 g of N-cyclopentyl-4-difluoromethoxybenzylamine, 20.2 g of triethylamine and 200 ml of dioxane. Under cooling with water 17.4 g of phenylisocyanide dichloride was added dropwise followed by stirring under reflux for 2 hours. After cooling the reaction mixture was treated in the same way as in Example 1 to afford 53.6 g of the title compound as pale brown crystals. The product was recrystallized from a hexane-ethyl acetate mixture to give white crystals (yield 28.2 g), m.p. 86.5-88 °C.

Preparative Example 3 Preparation of 1,3-dimethyl-1,3-di(4-chlorobenzyl)-2-(2,6-dimethylphenyl) guanidine (Compound No. 56)

In a 500-ml four-necked flask were placed 31.1 g of N-methyl-4-chlorobenzylamine and 200 ml of acetonitrile. After cooling with water 20.2 g of 2,6-dimethylphenylisocyanide dichloride was added dropwise followed by stirring at 50 °C for 3 hours. After cooling the reaction mixture was treated in the same way as in Example 1 to afford 37.4 g of the title compound as a pale yellow oil. The product was purified by column chromatography on silica gel using a hexane-acetone mixture to give a colorless oil (yield 30.3 g), $n_D^{23} = 1.5603$.

Preparative Example 4 Preparation of 1,3-dicyclohexyl-1,3-dibenzyl-2-(4-methylphenyl) guanidine (Compound No. 76)

In a 500-ml four-necked flask were placed 37.8 g of N-cyclohexylbenzylamine, 20.2 g of triethylamine and 200 ml of dioxane. Under cooling with water 18.8 g of 4-methylphenylisocyanide dichloride was added dropwise followed by stirring under reflux for 2 hours. After cooling, the reaction mixture was treated in the same way as in Example 1 to afford 25.6 g of the title compound as pale brown crystals. The product was recrystallized from a hexane-ethyl acetate mixture to give white crystals (yield 13.4 g), m.p. 128-130 °C.

Method for formulating fungicides for agriculture and horticulture

The agricultural and horticultural fungicides of the invention can be prepared by formulating the

compounds of the above-mentioned formula (I) in a conventional form. Thus, the compounds of the formula (I) may be compounded with suitable carriers and adjuvants, e.g., surface active agents, binders, stabilizers or the like to formulate wettable powder, emulsifiable concentrate, liquid formulation, sol (flowable formulation), oil solution, dust, DL (Driftless type) dust, microgranules, coarse dust or the like. The content (%) of the present compound in these formulations may range from 1 to 90% by weight for wettable powder, emulsifiable concentrate, liquid formulation, sol and oil solution, from 0.5 to 10% by weight for dust, DL dust, microgranules and coarse dust.

The method for using the agricultural and horticultural fungicides of the invention is illustrated below. In the application of wettable powder, liquid formulation, emulsifiable concentrate, sol (flowable formulation) and oil solution, they are diluted to 500-2000 times with water and generally adjusted to a solution containing 1 to 10000 ppm of the active ingredient. This diluted solution is sprayed over the foliage in the disease infection area of plant in an amount of 50 to 300 liters, usually 100 to 200 liters per 10 ares.

The liquid formulation, emulsifiable concentrate and sol (flowable formulation) are sprayed as a concentrated solution without dilution with water or a solution diluted to 10 times or less with water, principally as ultra low volume spray (LV spray, ULV spray) for aerial application in an amount of about 50 to 3000 ml per 10 ares using helicopter or the like.

The dust, DL dust, microgranules and coarse dust are applied to the foliage in the disease infection area of plant, in soil and onto the surface of soil or water, in an amount of 2 to 5 kg per 10 ares (about 50 to 500 g as active ingredient).

The method for formulating the inventive compounds of the general formula (I) into agricultural and horticultural fungicides will be illustrated below in Examples 5-8.

Example 5 Dust

A homogeneous mixture of 2 parts of Compound No. 13, 1 part of PAP (Modifier of physical properties) and 97 parts of clay was pulverized to obtain a dust containing 0.2% of the active ingredient.

Example 6 Wettable powder

A homogeneous mixture of 20 parts of Compound No. 21, 3 parts of potassium alkylbenzenesulfonate, 5 parts of polyoxyethylene nonyl phenyl ether and 72 parts of clay was pulverized to give a wettable powder containing 20% of the active ingredient.

Example 7 Emulsifiable concentrate

30 parts of Compound No. 3, 40 parts of methyl ethyl ketone and 30 parts of polyoxyethylene nonyl phenyl ether were mixed and dissolved to give an emulsifiable concentrate containing 30% of the active ingredient.

Example 8 Sol

40 parts of Compound No. 25, 2 parts of lauryl sulfate, 2 parts of sodium alkyl naphthalenesulfonate, 1 part of acetoxypyrrol cellulose and 55 parts of water were blended homogeneously to give a sol containing 40% of the active ingredient.

The novel compounds of the invention possess high control effects on mildew, blight, powdery mildew and rust which are serious diseases of fruit trees, vegetables or cereals, and are useful as agricultural and horticultural fungicides.

Usefulness and embodiments of the inventive compounds of the formula (I) will be shown with reference to Test Example 1-4.

Test Example 1 Test for the control effect on cucumber downy mildew

The second-leaf stage young seedlings of cucumber (variety: Sagami hanjiro) soil-cultured in a pot of 9 cm in diameter in a greenhouse was applied 20 ml per pot of a test liquid which was prepared by diluting a wettable powder prepared according to Example 6 to a predetermined concentration. One day after the application, the seedling was inoculated by spraying with a spore suspension of cucumber downy mildew fungus (*Pseudoperonospora cubensis*) which had been prepared by scraping spores with a wet brush off a fungus-infected leaf, suspending the spores in a 50 ppm aqueous solution of sticker (polyoxyethylene alkyl

ether) and adjusting the suspension to a concentration of 5×10^5 spores/ml. The seedlings were allowed to stand for 2 days at 20°C and 100% humidity to induce the disease development. Six days after the inoculation, the area (%) of lesions per leaf was observed, and the control value (%) was calculated according to the equation shown below.

The test was conducted in two-series of pots for each concentration of the test liquid, and an average control value (%) was calculated to make an evaluation on the basis of the index given below. Furthermore, the phytotoxicity against cucumber was assessed. The results are shown in Table 2.

It is to be noted that the same evaluation of the fungicidal effect and the same examination of the phytotoxicity score as in this example were run also in Test Examples 2-4.

$$\text{Control value (\%)} = \left(1 - \frac{\text{Area of lesions per leaf in treated pot}}{\text{Area of lesions per leaf in untreated pot}}\right) \times 100$$

<u>Index of fungicidal effect</u>	<u>Control value</u>
5	100%
4	80 - < 100%
3	60 - < 80%
2	40 - < 60%
1	20 - < 40%
0	< 20%
<u>Index of phytotoxicity</u>	
5: Very severe	
4: Severe	
3: Moderate	
2: Some	
1: Slight	
0: No	

Table 2

5	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
	1	100	4	0
10	2	100	4	0
	3	100	5	0
15	4	100	4	0
	5	100	5	0
	6	100	5	0
20	7	100	5	0
	8	100	5	0

25

30

35

40

45

50

55

Table 2 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	9	100	4	0
10	10	100	5	0
	11	100	4	0
15	12	100	5	0
	13	100	4	0
	14	100	5	0
20	15	100	5	0
	16	100	5	0
25	17	100	4	0
	18	100	5	0
	19	100	5	0
30	20	100	5	0
	21	100	5	0
35	22	100	4	0
	23	100	5	0
	24	100	5	0
40	25	100	5	0
	26	100	5	0
45	27	100	5	0
	28	100	4	0
	29	100	5	0
50	30	100	5	0

55

Table 2 (cont'd)

5	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
	31	100	4	0
10	32	100	5	0
	33	100	4	0
15	34	100	5	0
	35	100	5	0
	36	100	5	0
20	37	100	5	0
	38	100	5	0
25	39	100	5	0
	40	100	5	0
	41	100	5	0
30	42	100	5	0
	43	100	5	0
35	44	100	5	0
	45	100	5	0
	46	100	5	0
40	47	100	5	0
	48	100	5	0
45	49	100	4	0
	50	100	4	0
	51	100	4	0
50	52	100	4	0

55

Table 2 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	53	100	5	0
10	54	100	5	0
	55	100	5	0
15	56	100	5	0
	57	100	4	0
	58	100	4	0
20	59	100	4	0
	60	100	5	0
25	61	100	4	0
	62	100	4	0
	63	100	5	0
30	64	100	4	0
	65	100	5	0
35	66	100	5	0
	67	100	4	0
	68	100	4	0
40	69	100	4	0
	70	100	5	0
45	71	100	5	0
	72	100	4	0
	73	100	4	0
50	74	100	5	0

55

Table 2 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	75	100	5	0
10	76	100	5	0
	77	100	5	0
15	78	100	5	0
	79	100	4	0
	80	100	5	0
20	81	100	5	0
	82	100	5	0
25	83	100	4	0
	84	100	5	0
	85	100	5	0
30	86	100	5	0
	87	100	5	0
35	88	100	5	0
	89	100	4	0
	90	100	4	0
40	91	100	4	0
	92	100	4	0
45	93	100	5	0
	94	100	5	0
	95	100	5	0
50	96	100	5	0


55

Table 2 (cont'd)


Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
97	100	4	0
98	100	5	0
99	100	4	0
100	100	5	0
101	100	5	0
102	100	4	0
103	100	5	0
Comparative compound A	100	2	3
Comparative compound B	100	4	1
Untreated	-	0 (93.6)	-

(Note) The parenthesized number in the untreated indicates percent of lesion area per leaf.

Comparative compound A:



(Comparative compound B is described in Japanese Patent Publication No. 29742/1968 and No. 9846/1975)



(Generic name: Chlorothalonil)

Test Example 2 Test for control effect on tomato late blight

The young seedlings of tomato (variety: Toko K, the fifth leaf stage seedlings) soil cultured in a vinyl pot of 9 cm in diameter in a greenhouse was applied by means of an auto-sprayer 60 ml per 3 pots of a test liquid. The test liquid had been prepared by diluting a wettable powder prepared according to the procedure of Example 6 to a predetermined concentration. On the next day after completion of the spraying of the test liquid, the leaves of treated seedlings were inoculated by spraying a zoosporangium suspension of pathogenic fungus (*Phytophthora infestans*) by a hand sprayer. The zoosporangia had been rinsed out of the fungus cultivated on potato tubers at 20°C for 5 days, and the suspension prepared at a zoosporangim concentration of 10⁵/ml. The seedlings were then kept in a moist chamber at 20°C and 100% humidity for 5 days followed by observation of the area (%) of lesions on the first to fourth leaves. Percent average area of lesions per leaf was calculated, and control value (%) determined in comparison with the untreated plot.

The test was conducted in two-series of pots for each concentration of the test liquid, and average control value (%) determined, which was translated into evaluation score.

Furthermore, the phytotoxicity against the tomato seedling of the test compound was investigated according to the same procedure as in Test Example 1. The results obtained are as shown in Table 3.

$$\text{Control value (\%)} = \left(1 - \frac{\begin{array}{c} \% \text{ of lesions per leaf} \\ \text{in treated plot} \end{array}}{\begin{array}{c} \% \text{ of lesions per leaf} \\ \text{in untreated plot} \end{array}} \right) \times 100$$

Table 3

5	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
	1	100	4	0
10	2	100	5	0
	3	100	5	0
	4	100	4	0
15	5	100	4	0
	6	100	5	0
20	7	100	5	0
	8	100	5	0

25

30

35

40

45

50

55

Table 3 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	9	100	4	0
10	10	100	5	0
	11	100	4	0
15	12	100	5	0
	13	100	4	0
	14	100	5	0
20	15	100	5	0
	16	100	5	0
25	17	100	4	0
	18	100	5	0
	19	100	5	0
30	20	100	5	0
	21	100	5	0
35	22	100	4	0
	23	100	5	0
	24	100	5	0
40	25	100	5	0
	26	100	5	0
45	27	100	5	0
	28	100	5	0
	29	100	5	0
50	30	100	5	0

55

Table 3 (cont'd)

5	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
	31	100	5	0
10	32	100	4	0
	33	100	5	0
15	34	100	4	0
	35	100	4	0
	36	100	4	0
20	37	100	5	0
	38	100	4	0
25	39	100	5	0
	40	100	4	0
	41	100	4	0
30	42	100	4	0
	43	100	4	0
35	44	100	5	0
	45	100	5	0
	46	100	4	0
40	47	100	4	0
	48	100	5	0
45	49	100	4	0
	50	100	4	0
	51	100	4	0
50	52	100	5	0

55

Table 3 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	53	100	5	0
10	54	100	5	0
	55	100	5	0
	56	100	5	0
15	57	100	5	0
	58	100	4	0
20	59	100	4	0
	60	100	5	0
	61	100	5	0
25	62	100	4	0
	63	100	4	0
30	64	100	5	0
	65	100	5	0
	66	100	5	0
35	67	100	4	0
	68	100	4	0
40	69	100	4	0
	70	100	5	0
	71	100	5	0
45	72	100	5	0
	73	100	4	0
50	74	100	4	0

55

Table 3 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	75	100	5	0
10	76	100	5	0
	77	100	5	0
15	78	100	5	0
	79	100	4	0
	80	100	5	0
20	81	100	4	0
	82	100	5	0
25	83	100	4	0
	84	100	4	0
	85	100	4	0
30	86	100	4	0
	87	100	5	0
35	88	100	5	0
	89	100	5	0
	90	100	5	0
40	91	100	5	0
	92	100	4	0
45	93	100	5	0
	94	100	5	0
	95	100	5	0
50	96	100	4	0

55

Table 3 (cont'd)

Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
97	100	4	0
98	100	5	0
99	100	5	0
100	100	5	0
101	100	4	0
102	100	5	0
103	100	4	0
Comparative A	100	1	3
compound B	100	4	1
Untreated	-	0 (91.6)	-

Note 1) The comparative compounds A and B are the same as those shown in Table 2.

2) The parenthesized figure in the untreated indicates the number of lesion per leaf.

Test Example 3 Test for the control effect on barley powdery mildew

Over the first-leaf stage seedlings of barley (variety: Azumagolden) soil-cultured in a biscuit pot of 9 cm in diameter in a greenhouse were sprayed 10 ml of a test liquid which had been prepared by diluting a wettable powder prepared according to Example 6 to a predetermined concentration.

After allowing to stand overnight, the seedlings were inoculated by spraying with a spore suspension of powdery mildew fungus (*Erysiphe graminis*). Seven days after the inoculation, the number of lesion infected with barley powdery mildew was observed, and the control value (%) was calculated according to the formula below and translated into evaluation index. Furthermore, the phytotoxicity against the barley of the test compound was examined on the same ratings as described in Test Example 1. The results are shown in Table 4.

$$\text{Control value (\%)} = \left(1 - \frac{\text{Number of the lesion in treated pot}}{\text{Number of lesion in untreated pot}} \right) \times 100$$

Table 4

Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
1	250	5	0
2	250	4	0
3	250	5	0
4	250	5	0
5	250	5	0
6	250	5	0
7	250	5	0
8	250	5	0

Table 4 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	9	250	4	0
10	10	250	5	0
	11	250	4	0
15	12	250	5	0
	13	250	4	0
	14	250	5	0
20	15	250	5	0
	16	250	5	0
25	17	250	5	0
	18	250	5	0
	19	250	5	0
30	20	250	5	0
	21	250	5	0
35	22	250	5	0
	23	250	5	0
	24	250	5	0
40	25	250	5	0
	26	250	5	0
45	27	250	5	0
	28	250	5	0
	29	250	5	0
50	30	250	5	0

55

Table 4 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	31	250	5	0
10	32	250	5	0
	33	250	4	0
15	34	250	5	0
	35	250	5	0
	36	250	5	0
20	37	250	5	0
	38	250	5	0
25	39	250	5	0
	40	250	4	0
	41	250	4	0
30	42	250	4	0
	43	250	5	0
35	44	250	5	0
	45	250	4	0
	46	250	5	0
40	47	250	5	0
	48	250	4	0
45	49	250	4	0
	50	250	4	0
	51	250	4	0
50	52	250	5	0

55

Table 4 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	53	250	5	0
10	54	250	5	0
	55	250	5	0
	56	250	5	0
15	57	250	4	0
	58	250	4	0
20	59	250	4	0
	60	250	5	0
	61	250	4	0
25	62	250	4	0
	63	250	5	0
30	64	250	5	0
	65	250	5	0
	66	250	5	0
35	67	250	5	0
	68	250	5	0
40	69	250	4	0
	70	250	4	0
	71	250	4	0
45	72	250	4	0
	73	250	4	0
50	74	250	4	0

Table 4 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	75	250	5	0
10	76	250	5	0
	77	250	5	0
15	78	250	5	0
	79	250	5	0
	80	250	4	0
20	81	250	5	0
	82	250	4	0
25	83	250	4	0
	84	250	5	0
	85	250	5	0
30	86	250	5	0
	87	250	5	0
35	88	250	5	0
	89	250	4	0
	90	250	5	0
40	91	250	4	0
	92	250	4	0
45	93	250	5	0
	94	250	5	0
	95	250	5	0
50	96	250	5	0

55


Table 4 (cont'd)

Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
97	250	4	0
98	250	5	0
99	250	4	0
100	250	4	0
101	250	5	0
102	250	5	0
103	250	5	0
Comparative A	250	2	2
compound C	250	4	1
Untreated	-	0 (50.6)	-

Note 1) The comparative compound A is the same as the one shown in Table 2.

2) The parenthesized figure in the untreated indicates the number of the lesion per leaf.

Comparative compound C:



(Generic name: Quinomethionate)

Test Example 4 Test for the control effect on wheat leaf rust

Over the 1st-leaf stage young seedlings of wheat (variety: Norin No. 61) soil-cultured in a biscuit pot of 9 cm in diameter in a greenhouse was sprayed 20 ml per 3 pots of a test liquid which had been prepared by diluting a wettable powder prepared according to Example 6 to a predetermined concentration. One day after the spraying of the test liquid, the leaves to be treated were inoculated by spraying with a spore suspension of leaf rust fungus. The spore suspension had been prepared by suspending uredospores of wheat leaf rust fungus (Puccinia recondita) formed on the leaf of wheat in a distilled water containing 50

ppm of Tween 20® (trade name of polyoxyethylene sorbitan monolaurate, Kao Soap Co., Ltd.) so as to give about 50 spores as observed at one view under a microscope (x 150). After keeping overnight in a moist chamber at 20 °C, the treated seedlings were transferred to a disease development greenhouse at 20 °C to induce an incidence. Ten days after the inoculation, the seedlings were taken from the greenhouse and the number of uredosporus per leaf was counted and the control value (%) was calculated on the basis of the following formula. The test was performed in three-series of pots for each concentration of test compound to determine an average control value (%) which was translated into the evaluation index. Furthermore, the phytotoxicity against wheat of the test compound was examined on the same ratings as described in Test Example 1. The results are shown in Table 5.

$$\text{Control value (\%)} = \left(1 - \frac{\text{Number of uredosporus per treated leaf}}{\text{Number of uredosporus per untreated leaf}} \right) \times 100$$

Table 5

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	1	250	4	0
10	2	250	5	0
	3	250	5	0
	4	250	4	0
15	5	250	5	0
	6	250	5	0
20	7	250	5	0
	8	250	5	0
	9	250	5	0
25	10	250	5	0
	11	250	4	0
30	12	250	5	0
	13	250	4	0
	14	250	5	0
35	15	250	5	0
	16	250	5	0
40	17	250	4	0

45

50

55

Table 5 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	18	250	5	0
10	19	250	5	0
	20	250	5	0
	21	250	5	0
15	22	250	4	0
	23	250	5	0
20	24	250	5	0
	25	250	5	0
	26	250	5	0
25	27	250	5	0
	28	250	5	0
30	29	250	5	0
	30	250	5	0
	31	250	5	0
35	32	250	5	0
	33	250	4	0
40	34	250	5	0
	35	250	5	0
	36	250	4	0
45	37	250	5	0
	38	250	5	0
50	39	250	5	0

55

Table 5 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	40	250	4	0
10	41	250	5	0
	42	250	4	0
	43	250	4	0
15	44	250	5	0
	45	250	5	0
20	46	250	5	0
	47	250	5	0
	48	250	5	0
25	49	250	4	0
	50	250	4	0
30	51	250	4	0
	52	250	5	0
	53	250	5	0
35	54	250	5	0
	55	250	5	0
40	56	250	5	0
	57	250	5	0
	58	250	5	0
45	59	250	4	0
	60	250	5	0
50	61	250	4	0

55

Table 5 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5	62	250	5	0
10	63	250	5	0
	64	250	4	0
15	65	250	5	0
	66	250	5	0
	67	250	4	0
20	68	250	4	0
	69	250	5	0
25	70	250	5	0
	71	250	4	0
	72	250	4	0
30	73	250	5	0
	74	250	4	0
35	75	250	4	0
	76	250	5	0
	77	250	5	0
40	78	250	4	0
	79	250	4	0
45	80	250	5	0
	81	250	4	0
	82	250	5	0
50	83	250	4	0

55

Table 5 (cont'd)

	Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
5				
10	84	250	5	0
	85	250	5	0
	86	250	4	0
15	87	250	5	0
	88	250	5	0
20	89	250	4	0
	90	250	5	0
	91	250	4	0
25	92	250	4	0
	93	250	5	0
30	94	250	4	0
	95	250	5	0
	96	250	4	0
35	97	250	4	0
	98	250	5	0
40	99	250	5	0
	100	250	4	0
	101	250	5	0
45	102	250	4	0
	103	250	5	0
50				
55				


Table 5 (cont'd)

Test compound No.	Concentration (ppm)	Index of the effect	Phytotoxicity
Comparative A	250	1	2
compound D	250	4	1
untreated	—	0 (49.2)	—

Note 1) The comparative compound A is the same as that shown in Table.2.


2) The parenthesized number in the untreated indicates the number of the lesion per leaf.

Comparative compound D:



Claims

1. A guanidine derivative of formula (I)



wherein R is hydrogen, (C₁-C₁₀)alkyl, (C₂-C₆)alkenyl, (C₂-C₄)-alkynyl, (C₅-C₆)cycloalkyl(C₁-C₆)alkyl or (C₅-C₇)cycloalkyl optionally substituted with (C₁-C₆)alkyl, X is hydrogen, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy, Y is hydrogen, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy, R¹ and R² are the same or different and each is hydrogen or methyl, and l, m and n are the same or different and each is an integer of 1 or 2 provided that when R, X, Y, R¹ and R² are all hydrogen, l is 2, or an acid addition salt

thereof.

2. A compound according to claim 1 wherein R is (C₅-C₆)-cycloalkyl, X is halogen, (C₁-C₆)alkyl or (C₁-C₆)-alkoxy, Y is hydrogen, R¹ and R² are both hydrogen and l, m and n are all 1, or an acid addition salt thereof.


3. An acid addition salt according to claim 1 or claim 2 which is a salt of an inorganic acid or a carboxylic or sulphonic acid.

4. An acid addition salt according to claim 3 which is a salt of hydrochloric, sulphuric, nitric, phosphoric, acetic, propionic, glycolic, lactic, citric, or benzene sulphonic acid.

5. A fungicide for agriculture and horticulture comprising, as active ingredient, a dibenzylguanidine derivative according to any one of claims 1 to 4.


6. Use of a compound of formula (I) according to any one of claims 1 to 4 as an agricultural or horticultural fungicide.

7. A process for producing a compound of formula (I) according to any one of claims 1 to 4 comprising reacting a compound of formula (II).



(II)

wherein Y and n are as defined in relation to the compound of formula (I) with a compound of formula (III)



wherein R, X, R¹, R², l and m are as defined in relation to the compound of formula (I) in the presence of a base.

8. A process according to claim 7 wherein the compound of formula (III) is used in excess as the base.



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 429 281 A3**

EUROPEAN PATENT APPLICATION

Application number: **90312579.7**

Int. Cl.⁵: **C07C 279/18, A01N 47/44**

Date of filing: **19.11.90**

Priority: **20.11.89 JP 299655/89**
04.10.90 JP 265230/90

Date of publication of application:
29.05.91 Bulletin 91/22

Designated Contracting States:
DE FR GB IT

Date of deferred publication of the search report:
27.05.92 Bulletin 92/22

Applicant: **HOKKO CHEMICAL INDUSTRY CO., LTD**
4-20, 4-chome, Nihonbashi-Hongoku-cho
Chuo-ku
Tokyo(JP)


Inventor: **Ishikawa, Hiromichi**
40-12, Asahi-cho 5-chome
Atsugi-shi, Kanagawa-ken(JP)

Inventor: **Umeda, Takashi**
5-2-306, Sagamidaidanchi
Sagamihara-shi, Kanagawa-ken(JP)
Inventor: **Onoue, Shinji c/o Hokko Kagaku-Ryo**
2385, Toda
Atsugi-shi, Kanagawa-ken(JP)
Inventor: **Kajikawa, Kazuo**
c/o Hokko Kagaku-Ryo, 2385, Toda
Atsugi-shi, Kanagawa-ken(JP)
Inventor: **Shibata, Toshihiro**
c/o Hokko Kagaku-Ryo, 2385, Toda
Atsugi-shi, Kanagawa-ken(JP)
Inventor: **Ohshima, Hiroshi**
B-22-19, 36-20, Tsutsumi
Chigasaki-shi, Kanagawa-ken(JP)

Representative: **Cresswell, Thomas Anthony et al**
J.A. Kemp & Co. 14 South Square Gray's Inn
London WC1R 5LX(GB)

Guanidine derivatives and fungicides for agriculture and horticulture containing the same.

The guanidines of formula (I)



wherein R is hydrogen, (C₁-C₁₀)alkyl, (C₂-C₆)alkenyl, (C₂-C₄)-alkynyl, (C₅-C₆)cycloalkyl, (C₁-C₆)alkyl or (C₅-C₇)cycloalkyl optionally substituted with (C₁-C₆)alkyl, X is hydrogen, halogen, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, halo(C₁-C₆)alkyl or halo(C₁-C₆)alkoxy, Y is hydrogen, halogen, (C₁-C₆)alkyl or (C₁-C₆)alkoxy, R¹ and R² are the same or different and each is hydrogen or methyl, and l, m and n are the same or different and each is an integer of 1 or 2 provided that when R, X, Y, R¹ and R² are all hydrogen, l is 2, and acid addition salts thereof possess high control effects on mildew, blight, powdery mildew and rust which are serious diseases of fruit trees, vegetables or cereals, and are useful as agricultural and horticultural fungicides.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 31 2579

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	CHEMICAL ABSTRACTS, vol. 83, no. 15, 13 October 1975, Columbus, Ohio, US; abstract no. 127548D, page 164 ;column 2 ; in combination with formula index volume 83 page 1870f column 3 lines 24 - 26 * abstract *	1,6	C07C279/18 A01N47/44
D,X	& JP-A-5 009 846 (KUMIAI CHEMICAL INDUSTRY CO., LTD.) 16 April 1975	1,6	
A	DE-A-3 108 564 (BAYER AG) * claims 1,5 *	1,7	
A	US-A-3 976 643 (J. DIAMOND ET AL) * formula I column 1 *	1	
A	US-A-3 320 229 (K. SZABO ET AL) * column 2 *	1,7	
			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
			C07C A01N
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 19 MARCH 1992	Examiner KAPTEYN H.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons * : member of the same patent family, corresponding document	

EPO FORM 1503 03.82 (P0401)